

Evaluation of antioxidant and antifungal activities of several plants against agents of postharvest citrus sour rot and green mould rot

Received for publication, July, 7, 2016

Accepted, March, 21, 2017

BİLAL BALKAN^{1*}, HALİDE AYDOĞDU², SEDA BALKAN³, BUKET AŞKIN⁴, HÜSEYİN ERSOY⁵

^{1*} Vocational School of Technical Sciences, Kırklareli University, Kırklareli, Turkey

² Arda Vocational College, Trakya University, Edirne, Turkey

³ Science and Art Faculty, Department of Molecular Biology and Genetic, Kırklareli University, Kırklareli, Turkey

⁴ Engineering Faculty, Department of Food Engineering, Kırklareli University, Kırklareli, Turkey

⁵ Science Faculty, Department of Biology, Trakya University, Edirne, Turkey

*Address for correspondence to: bilalbalkan@hotmail.com

Abstract

The antifungal activities of chloroform extracts of 10 plants species belonging to Lamiaceae family, which were collected from Kırklareli (Turkey), against *Geotrichum candidum*, the agent of postharvest citrus sour rot and *Penicillium digitatum*, the agent of postharvest citrus green mould rot, were researched. The lowest Minimum Inhibitory Concentration (MIC) values against *G. candidum* and *P. digitatum* were obtained in the extract of *Marrubium peregrinum* L. (250 and 125 µg/ml). In 1000 µg/ml, the extracts of *Melissa officinalis* showed 100% inhibition on the spore germination of *G. candidum* and *P. digitatum*. In the Scanning Electron Microscope (SEM) observations of *G. candidum* and *P. digitatum* that was subjected to *M. peregrinum* extract (4MIC) degenerative changes in the hyphal morphology were seen in the form of cell wall degradation, lysis and collapsing. The highest values of total phenolics were obtained from *Mentha pulegium* extracts (739.57 mg GAE/g). The lowest EC₅₀ values (0.08 mg/ml) were found in the extracts of *M. peregrinum* and *Sideritis montana*. The highest flavanol content was determined from *M. officinalis* extracts (12.71 mg CE/mg). This study demonstrates *M. peregrinum* extracts may possess high antifungal activity against *G. candidum* and *P. digitatum*.

Keywords: *Geotrichum candidum*, *Penicillium digitatum*, Antifungal activity, Plant extract, SEM

1. Introduction

The wounds occurring on the fruits and vegetables during harvest, transportation and/or processing may be infected with pathogenic fungi during storage to cause microbial decay, the primary problem for the food industry (AMEZIANE & al. [1]; PHILLIPS & al. [2]). Due to the infections the fungi caused on the foods, the nutritional value of the food decrease and they become unfit for consuming. Furthermore, the lipid peroxidation caused by the free radical oxidation also damages the foods greatly (PRAKASH & al. [3]). Additionally, the fact that free radicals cause degenerative diseases has been proven by many studies (CHEN & al. [4]). Turkey is an important growing ground for citrus fruits, producing more than 3.5 million tons of citrus fruits annually, and holds 9th place in the world citrus production (YEŞİLOĞLU & al. [5]). Due to their low pH, high water concentration and nutritional components citrus are very sensitive to attacks from pathogenic fungi in the period between harvest and consumption (TRIPATHI & al. [6]). The most common and severe diseases in citrus worldwide are sour rot caused by *Geotrichum candidum* (Link ex Pers) and green mould rot caused by *Penicillium digitatum* (Pers.:Fr.). The decay in fruits and vegetables after harvest can be reduced by 90-95% using fungicides. However, the losses may increase by 50%

without fungicide treatment (PHILLIPS & al. [2]). The green mould rot in fruits and vegetables after harvest may be controlled by intensive usage of chemical fungicide pre and post-harvest in many countries including Turkey (SOYLU & al. [7]). For decades, benomyl, thiabendazole, imazalil, and sodium ortho-phenylphenate (SOPP), which belong to the benzimidazole (BZI), sterol demethylation inhibitor (DMI), and phenyl-phenol chemical classes, respectively, were the most commonly used fungicides until the recent introduction to the market of pyrimethanil (anilinopyrimidine) and fludioxonil (phenylpyrrole) (AMIRI AND BOMPEIX, [8]). Sour rot is only controlled by Guazatine, a chemical not authorized in several countries, and not by imazalil and thiabendazole, mainly used in post-harvest chemical treatments (TALIBI & al. [9]). The usage of these fungicides are restricted because of their high and acute toxicity, long degradation periods, residues in the food chain, and the chronic poisoning through continuous intake in small quantities (SHUKLA & al. [10]). However, the increased resistance of fungi against synthetic chemicals causes problems in the effective uses of fungicides (AL-REZA & al. [11]). In addition to the loss of activities of the fungicides, their negative environmental effects force people (PASSONE & al. [12]) to discover new crop protection methods and new types of selective control alternatives that does minimal damage to the environment and human health (SOYLU & al. [13]). It is essential to optimize alternative methods that have minimal environmental and human damage and have a different effect mechanism in order to avoid the fungicide resistance in microorganisms (PASSONE & al. [12]). Therefore, researchers are screening the plant extracts for an alternative method (ABDEL-MONAIM & al. [14]). Plants are seen as the source for effective chemical materials and provide reversible sources for harmless fungicides. A typical characteristic of plants is that they produce nitrogen free and nitrogen containing secondary metabolites. In most cases, these secondary metabolical products act as a defensive mechanism against microorganisms, insects and herbivores. Such products from higher plants are relatively broad-spectrum, bio-efficacious, economical, and presumably safer for humans and the environment and can be ideal candidates for use as agrochemicals (PASSONE & al. [15]). Meanwhile, plants act as singlet and triplet oxygen quenchers, free radical scavengers and enzyme inhibitors. These protective biological effects are attributed to the polyphenol contents of plants (CHEN & al. [4]). The systematic screening of the interaction between microorganisms and plant products is one of the main ways in researching the biological active substances of plants. Plant products whose antimicrobial spectrum has been recognized can be used as antimicrobial compounds or as an adjuvant to other antimicrobial compounds in order to improve their action (DE SOUZA& al. [16]). The goals of this study were; (I) The testing of the antifungal activities of chloroform extracts of 10 plants species belonging to Lamiaceae family which were collected from Kırklareli (Turkey) against *G. candidum* and *P. digitatum* using Clinic and Laboratory Standards Institute (CLSI, [17]) procedures, (II) Examining the effects on spore germination and hyphal morphology of these fungi, (III) The analyzing of the total phenolic, flavanol contents, antiradical activities and antioxidant capacities.

2. Materials and Methods

Plant materials

Ten plant materials were collected from different locations in Kırklareli (Turkey) in the months of June and July 2014. All of the samples were identified by Dr. Hüseyin ERSOY and were deposited in the herbarium laboratory in faculty of sciences, Trakya University, Edirne, Turkey. Plant samples were cleaned and dried in the shade, then grinded to a fine powder using a laboratory grinding mill (A11B, IKA) and stored in the dark at 4 °C until use.

Pathogens

Cultures of the fungal pathogens *Geotrichum candidum* and *Penicillium digitatum* used in the study were isolated from naturally decayed fruits. A pure culture of these fungi were maintained on potato dextrose agar (PDA) and stored at 4 °C and sub-cultured once a month.

Preparation of plant extracts

The powdered plant materials were separately extracted thrice at room temperature with chloroform solvent (500ml/100 g of plant material each run). The final extracts of each plant part were filtered using filter paper (Whatman) and were evaporated under vacuum at 40 °C using rotary vacuum evaporator. All extracts were stored in sterile glass bottles at room temperature.

Determination of minimum inhibitory concentration (MIC)

The pathogen inoculum consisted of aqueous spore suspension obtained from 7-days-old culture incubated at 25 °C. Spores were harvested by flooding with 5 ml of sterile distilled water containing 0.05% (v/v) Tween 80, and passing the suspension through two layers of sterile cheesecloth to remove hyphal fragments. The spore concentration was determined with the aid of a hemacytometer and adjusted to 10⁶ spores per ml with sterile distilled water. Using as spectrophotometer optic density (OD) and transmittance (T%) of the suspension at 530 nm were recorded. The spectrophotometric values of the inoculum suspension were stabilized for further studies. The 10⁶ spore/ml suspension was later watered down to 1:50 ratio with sterile distilled water. Meanwhile the 1:100 rationed suspensions was added to the PDA plate by the amount of 0.01 ml to confirm 10⁴ CFU/ml.

MIC of each extract was determined by using broth microdilution techniques according to the guidelines for filamentous fungi (M 38 A) (CLSI, [17]). MIC values were determined in RPMI-1640 (Sigma, St. Louis, MO, USA) buffered to pH 7.0 with MOPS. Microtiter trays were incubated at 25 °C. MICs were visually recorded at 96 h. For the assay, 40 µL of a stock solution of each extract in DMSO (50 mg/ml) was diluted with 960 µL of RPMI-1640 giving a solution of 2 mg/ml. 200 µL of this new solution was poured into the first well and then, 100 µL were transferred to the next well containing 100 µL of RPMI-1640. The same procedure was performed for all wells of the same file obtaining twofold dilutions of the extract. A volume of 100 µL of inoculum suspension was added to each well (with the exception of the sterility control where sterile water was added to the well instead) giving extract's concentrations from 1000 to 0.98 µg/ml and a final DMSO concentration ≤1%. End-points were defined as the lowest concentration of extract resulting in total inhibition (MIC) of visual growth compared to the growth in the control wells containing no antifungal (SVATEZ & al. [18]). Tests were carried out in duplicate.

Effect of plant extracts on spore germination

The germination of spores of *G. candidum* and *P. digitatum* was determined in concentrations of 125, 250, 500 and 1000 µg/ml of chloroform extract for the plants. Aliquots (40 µL) of a spore suspension (10⁶ spores/ml) were aseptically transferred in triplicate to sterile depression slides containing 40 µL with different concentrations of chloroform extracts, and then incubated at 25 °C for 96 h. Spore germination was estimated under a microscope using a micrometer. At least 100 spores of each replicate were observed. A spore was scored as germinated if the germ tube length was equal or exceed that of the spore (ASKARNE & al. [19]; ASKARNE & al. [20]). The data were expressed as percent spore germination inhibition and calculated by using the following formula:

GI (%) = $[(G_c - G_t)/G_c] \times 100$, where G_c and G_t represent the mean number of germinated spores in control and treated slides, respectively. Each treatment included three replicates and the experiment was conducted twice.

Effect of plant extract on hyphal morphology

For the determination of the effect of plant extracts on hyphal morphology, a mycelial agar disc from a 7-day old culture was first placed in the centre of PDA plate and incubated at 25 °C for 2 days to allow mycelium to grow into the medium. After 2 days of pre-incubation, different concentrations (4MIC) of plant extract used in vitro studies were dropped (onto covers of Petri dishes) and incubated at 25 °C for 3 days (SVATEZ & al. [18]). For SEM analysis, mycelial discs (1 cm in diameter) were fixed with 2.5% glutaraldehyde in 0.1 M phosphate-buffer (pH=7.2) for 2 h at room temperature. They were washed twice, each time for 10 min, in the same buffer. After fixation, the samples were dehydrated in a graded ethanol series (70%, 80%, 90%, and three times at 100%) for a period of 30 min in each series. The samples were examined and digital images captured using SEM (Quanta FEG 250) at an accelerating voltage of 5 kV.

Determination of total phenolic and flavanol content

Spectrophotometric measurements were performed by spectrophotometer (Shimadzu, UV mini-1240). Total phenolic contents (TPC) of the extracts were determined by the Folin–Ciocalteu colorimetric method (SINGLETON AND ROSSI, [21]). The estimation of phenolic compounds in the extracts was carried out in five replications and calculated by a calibration curve obtained with Gallic acid. Total phenolics were expressed as Gallic acid equivalents (mg GAE/g).

Total flavanol contents (TFC) were assayed colorimetrically by the DMAC method (ARNOUS & al. [22]). The absorbance of the extracts was measured at 640 nm and TFC in the extracts were expressed as catechin equivalents (mg CE/g).

Determination of antiradical activity

Different concentrations of extracts in methanol were added to 1 ml methanolic solution of DPPH (0.2 mM) in volumes of 1 ml. The mixture was shaken vigorously and allowed to stand at room temperature for 30 min. Then the absorbance was measured at 517 nm against methanol as the blank in a spectrophotometer (BAYDAR AND BAYDAR, [23]). DPPH radical free scavenging activity (%) was calculated according to the formula:

Antiradical activity (%) = $[(\text{Absorbance of control} - \text{Absorbance of sample}) / \text{Absorbance of control}] \times 100$

Extract concentrations providing 50% inhibition (EC_{50}) was calculated from the graph plotting inhibition percentage against extract concentrations. Gallic acid was used as a positive test control.

Statistical analysis

All data were subjected to statistical analysis of variance (ANOVA) using STATISTICA software and Tukey's multiple comparison tests. For these procedures, "SPSS for Windows, v 15.0" statistical package program was used and the values lower than $p < 0.05$ were considered significant.

3. Results and Discussion

Determination of minimum inhibitory concentration (MIC)

One of the technical variables in MIC studies is the amount of inoculums. The colour, shape and size of the spores can affect the spectrometric readings. The idea of fungal species having different optic density values for each genus is documented in M-38A. This is why the OD of inoculum amount must be standardized for each species. The OD and T% values of *G.*

candidum and *P. digitatum* inoculum suspensions (10^6 spores/ml) at 530 nm were standardized at 0.039, 91.3% and 0.017, 96%, respectively (Table 1).

Table 1. Optical Density and Transmittance (%) values of the spore suspension (10^6 spores/ml) of *Geotrichum candidum* and *Penicillium digitatum* at 530 nm.

Fungi	Optical Density	Transmittance (%)
<i>G. candidum</i>	0.039	91.3
<i>P. digitatum</i>	0.017	96

Chloroform extracts of all plants were evaluated for antifungal properties with the microbroth dilution method. It was reported that the plant chloroform extracts showed effective results in antifungal studies (ASHRAF & al. [24]; VOGT & al. [25]). A plant with MICs ≤ 1000 $\mu\text{g/ml}$ was considered active. Except *Lamium amplexicaule*, other plant extracts showed antifungal activity against tested fungi. In all of the tested plant species the MIC values were the lowest in *M. peregrinum* against *G. candidum* and *P. digitatum* (250 and 125 $\mu\text{g/ml}$), something which indicates this plant has more antifungal substances against *G. candidum* and *P. digitatum* than other plants (Table 2). The antifungal compound can lead to the inhibition of the fungus by suppressing the early stages of mycelium growth. Hydrophobicity is an important characteristic of the plant extracts and their components, which renders them more permeability and disturb cell structures by enabling partition of the lipid on the mitochondria and the cell membrane (RASTOGI AND MEHROTRA, [26]). The death of the cell can be from the extensive leakage from the living cells or the exit of the critical molecules and ions (DERESSA AND WAKJIRA, [27]).

Table 2. Antifungal activities (MIC in $\mu\text{g/ml}$) of the some plant chloroform extracts investigated against *Geotrichum candidum* and *Penicillium digitatum*.

Plant species	<i>G. candidum</i>	<i>P. digitatum</i>
<i>Ajuga laxmannii</i> (Murray) Benth.	1000	1000
<i>Lamium amplexicaule</i> L. var. <i>amplexicaule</i>	i	i
<i>Lamium maculatum</i> L.var. <i>maculatum</i>	500	500
<i>Marrubium peregrinum</i> L.	250	125
<i>Melissa officinalis</i> L. subsp. <i>officinalis</i>	500	1000
<i>Mentha pulegium</i> L.	500	1000
<i>Salvia verticillata</i> L.subsp. <i>verticillata</i>	1000	500
<i>Sideritis montana</i> L. subsp. <i>montana</i>	1000	1000
<i>Teucrium chamaedrys</i> L.subsp. <i>chamaedrys</i>	500	1000
<i>Teucrium polium</i> L. subsp. <i>polium</i>	500	500

i; inactive (MIC > 1000 $\mu\text{g/ml}$)

Effects of plant extracts on spore germination

The chloroform extracts of *L. maculatum*, *M. peregrinum*, *M. officinalis* and *M. pulegium* have high fungistatic effect (100% inhibition) against the spore germination of *G. candidum* at 1000 $\mu\text{g/ml}$. At 500 $\mu\text{g/ml}$, the chloroform extracts of *M. peregrinum* and *M. officinalis* inhibited the spore germination by 94.66% and 86%, respectively (Table 3).

Table 3. Effect of chloroform extracts on spore germination of *Geotrichum candidum*.

Plant species	Spore germination inhibition (%) ^a			
	Concentration of chloroform extracts ($\mu\text{g/ml}$)			
	125	250	500	1000
<i>Ajuga laxmannii</i>	0 ^a	0 ^a	0 ^a	50 ^b

<i>Lamium maculatum</i>	0 ^a	0 ^a	20.66 ^c	100 ^d
<i>Marrubium peregrinum</i>	0 ^a	4 ^e	94.66 ^{fd}	100 ^d
<i>Melissa officinalis</i>	0 ^a	0 ^a	86 ^f	100 ^d
<i>Mentha pulegium</i>	0 ^a	0 ^a	43.66 ^b	100 ^d
<i>Salvia verticillata</i>	0 ^a	0 ^a	0 ^a	59 ^b
<i>Sideritis montana</i>	0 ^a	0 ^a	0 ^a	91 ^f
<i>Teucrium chamaedrys</i>	0 ^a	0 ^a	0 ^a	81.66 ^f
<i>Teucrium polium</i>	0 ^a	0 ^a	12.66 ^c	62 ^b
Control	0 ^a	0 ^a	0 ^a	0 ^a

^a Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at P < 0.05 according to Tukey's multiple comparison tests.

At 1000 µg/ml, the chloroform extract of *M. officinalis* inhibited the spore germination of *P. digitatum* by 100%. *L. maculatum*, *M. peregrinum* and *M. pulegium* extracts showed 96.83%, 94% and 83% inhibition, respectively (Table 4). The antifungal activity of the metabolites produced by the plant can show differences based on the stages of fungal development (CARVALHO & al. [28]). Some plant metabolites act upon the mycelium growth of the fungus while some are effective on the spore germination (TALIBI & al. [9]; ASKARNE & al. [20]; GARDUNO-PIZANA & al. [29]). Cell wall altering its permeability or inhibiting the spore germination may be the general effects of the antifungal compounds acting (AMSALU & al. [30]).

Table 4. Effect of chloroform extracts on spore germination of *Penicillium digitatum*.

Plant species	Spore germination inhibition (%) ^a			
	Concentration of chloroform extracts (µg/ml)			
	125	250	500	1000
<i>Ajuga laxmannii</i>	0 ^a	0 ^a	0 ^a	10.33 ^b
<i>Lamium maculatum</i>	0 ^a	0 ^a	56.88 ^c	96.83 ^d
<i>Marrubium peregrinum</i>	21.66 ^b	26.66 ^{cb}	80.66 ^f	94 ^d
<i>Melissa officinalis</i>	0 ^a	9 ^b	85.33 ^f	100 ^d
<i>Mentha pulegium</i>	0 ^a	0 ^a	29.66 ^c	83 ^f
<i>Salvia verticillata</i>	0 ^a	0 ^a	0 ^a	6 ^b
<i>Sideritis montana</i>	0 ^a	0 ^a	0 ^a	35.33 ^e
<i>Teucrium chamaedrys</i>	0 ^a	0 ^a	0 ^a	22 ^b
<i>Teucrium polium</i>	5.33 ^b	7 ^b	8 ^b	9.66 ^b
Control	0 ^a	0 ^a	0 ^a	0 ^a

^a Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at P < 0.05 according to Tukey's multiple comparison tests.

Effects of extract on hyphal morphology

Important morphological differences were found when comparing the control to the hyphal morphologies of *G. candidum* and *P. digitatum* that have been treated with the 4MIC extract of *M. peregrinum* in PDA plates. When compared to hyphal structures that have a linear, orderly and homogenous cell wall, the SEM observations showed that the chloroform extract of *M. peregrinum* cause lysis, collapsing and cell wall degradation in *G. candidum*; cell wall degradation and collapsing in *P. digitatum* (Fig. 1 and Fig. 2). These results confirm that the chloroform extract of *M. peregrinum* is phytotoxic. The degradations in the cell wall may be related to the chloroform extract of *M. peregrinum* reacting with the enzymes regulating the cell wall, which in turn can destroy the fungal morphology and inhibit hyphal growth. Similar results were obtained by SOYLU & al. [13] at recording the effect of essential oil against cell wall growth of *Botrytis cinerea*.




Fig. 1. Scanning electron microscopy of *G. candidum* hyphae exposed to *M. peregrinum* chloroform extract (1000 µg/ml). (A and B) Healthy hyphae in control petri plates. (C and D) Effects of the extract on hyphal morphology. Note alterations in hyphal morphology including cell wall degradation (1.arrow), lysis (2.arrow), collapsing (3.arrow) (C) and cell wall degradation (1.arrow), lysis (2.arrow) (D).




Fig. 2. Scanning electron microscopy of *P. digitatum* hyphae exposed to *M. peregrinum* chloroform extract (500 µg/ml). (A and B) Healthy hyphae in control petri plates. (C and D) Effects of the extract on hyphal morphology. Note alterations in hyphal morphology including cell wall degradation (1.arrows), collapsing (2.arrow) (C) and cell wall degradation (1.arrows) (D).

Total phenolic and flavanoid contents

All plants make phenolic substances as secondary metabolites, whose roles are not known enough. This is why there are phenolic substances in varying amounts and quality in plant base foods. Phenolic substances are one of the biggest secondary metabolite groups which act as antimicrobial and antioxidant. The hydroxyl group's position and amount determine the antimicrobial properties, while the ability to bond with metals and scavenge free radicals determine the antioxidant properties of phenol groups. It has been suggested that phenolics have many effect types in preventing the growth of pathogenic agents. Because of this effects the enzymatic processes in energy production gets destroyed, the permeable membrane of the cell gets damaged or weakened, the physicochemical structure of the cell changes or nucleic acid synthesis gets affected (CUTTER, [31]). These bioactive compounds, interfere with the life process of the fungi, whether singly or in combination, by changing the physiological status of the cells, weakening or destroying the permeability barrier of the cell membrane, altering structural component synthesis, and acting as chelating agents (RONGAI & al. [32]). The highest TPC was found in *M. pulegium* (739.57 mg GA/g) while the lowest TPC was determined to be in *L. amplexicula* (87.51 mg GA/g) (Table 5). The TPC amounts found in the extracts of *S. verticillate* in methanol (TOSUN & al. [33]); *S. montana* in methanol, acetone and ethyl acetate (RADOJEVIĆ & al. [34]); *T. chamaedrys* in methanol, acetone, ethyl acetate and petrol ether (STANKOVIC & al. [35]); *T. polium* in methanol (STANKOVIC & al. [36]) were reported as 50.3, 97.85, 84.55, 49.05, 168.46, 169.50, 143.95, 30.39, 42.64 and 157.84 mg GA/g. These results are different from what we found. The reason for this may be because of the difference in the solvents used in extractions as well as the difference of the regions and the geographical conditions where the plants grew (TRIGUI & al. [37]).

The antioxidant activities of the extract were evaluated by the TFC as well. Colourless compounds called flavanols are the intermediate products in flavonoid biosynthesis. Flavanols easily react with the oxygen in the air by both chemical and enzymatic reactions. After the reactions they form proanthocyanidin by condensing. Proanthocyanidins are powerful antioxidants, they reduce free radicals. The highest flavanol content (12.71 mg CE/g) was found in the extract of *M. officinalis* (Table 5).

Table 5. Total phenolic, total flavanol contents and EC₅₀ values of chloroform extracts.

Plant species	Total phenolic (mg GA/g)*	Total flavanol (mg CE/g)*	EC ₅₀ (mg/ml)*
<i>Ajuga laxmannii</i>	150.23 ^a	5.73 ^a	58.20 ^a
<i>Lamium amplexicaule</i>	87.51 ^b	4.29 ^b	19.18 ^b
<i>Lamium maculatum</i>	377.90 ^c	8.32 ^c	0.32 ^c
<i>Marrubium peregrinum</i>	504.90 ^d	3.78 ^d	0.08 ^c
<i>Melissa officinalis</i>	379.90 ^c	12.71 ^e	2.28 ^c
<i>Mentha pulegium</i>	739.57 ^e	3.43 ^{fd}	0.15 ^c
<i>Salvia verticillata</i>	352.23 ^c	3.73 ^d	0.11 ^c
<i>Sideritis montana</i>	501.57 ^d	3.97 ^d	0.08 ^c
<i>Teucrium chamaedrys</i>	511.90 ^d	5.76 ^a	2.81 ^c
<i>Teucrium polium</i>	519.57 ^d	3.95 ^d	3.08 ^c
Gallic acid			1.86 ^c

* Each value represents the mean of three replicates. Means followed by different letters in each column are significantly different at P < 0.05.

DPPH free radical scavenging activity

The antioxidant activity of the extracts can also be evaluated by the determination of the EC₅₀ values corresponding to the amount of extract required to scavenge 50% of DPPH radicals

present in the reaction mixture. High EC₅₀ values indicate low antioxidant activity (BAYDAR AND BAYDAR, [23]). EC₅₀ values of *L. maculatum*, *M. peregrinum*, *M. pulegium*, *S. verticillate* and *S. montana* were less than the EC₅₀ value obtained for Gallic acid used as positive control (Table 5). Among the extracts, *M. peregrinum* and *S. montana* extracts were the most potent radical scavengers. STANKOVIC & al. [35] reported EC₅₀ values of 0.56 and 1.16 mg/ml in petrol ether extracts of *T. chamaedrys* leaves and flowers, respectively; RADOJEVIĆ & al. [34] reported EC₅₀ values of 0.031, 0.064 and 0.52 for methanol, acetone and ethyl acetate extracts of *S. montana*, respectively. The idea of the plant extract that has more total phenolics also shows more antiradical activity (BAYDAR AND BAYDAR, [23]) is shown again as in our study the high correlation (r=0.8981; p<0.01) between total phenolic amount and antioxidant capacities can be seen.

4. Conclusion

A. laxmannii, *L. amplexicaule*, *L. maculatum*, *M. peregrinum*, *M. officinalis*, *M. pulegium*, *S. verticillate*, *S. montana*, *T. chamaedrys* and *T. polium* species belonging to Lamiaceae family which can be found abundantly and grown easily in Kırklareli (Turkey), were tested with microdilution method against the primary fungi pathogen of citrus postharvest, *G. candidum* and *P. digitatum*. We can say that due to its high antifungal activity against *G. candidum* and *P. digitatum*, the chloroform extract of *M. peregrinum* has a high potential to be used as an alternative natural control agent instead of chemical fungicides. Also, among the tested plants, *L. maculatum*, *M. peregrinum*, *M. pulegium*, *S. verticillate* and *S. montana* have a lower EC₅₀ value than Gallic acid which shows that they have high antioxidant activity.

Acknowledgements

This study was supported financially by the Research Foundation of the University of Kırklareli, Turkey. Project number: KLUBAP/020.

References

1. N. AMEZIANE, H. BOUBAKER, H. BOUDYACH, F. MSANDA, A. JILAL, A.A. BENAOMAR. Antifungal Activity of Moroccan Plants Against Citrus Fruit Pathogens, *Agron. Sustain. Dev.*, 27: 273-277 (2007).
2. C.A. PHILLIPS, K. LAIRD, S.C. ALLEN. The Use of Citri-V™[®]-An Antimicrobial Citrus Essential Oil Vapour for The Control of *Penicillium chrysogenum*, *Aspergillus niger* and *Alternaria alternata* In vitro and on Food, *Food Research International*, 47: 310-314 (2012).
3. B. PRAKASH, P. SINGH, A. KEDIA, N.K. DUBEY. Assessment of Some Essential Oils as Food Preservatives Based on Antifungal, Antiaflatoxin, Antioxidant Activities and *In vivo* Efficacy in Food System, *Food Research International*, 49: 201-208 (2012).
4. F. CHEN, X. LONG, M. YU, Z. LIU, L. LIU. Phenolics and antifungal activities analysis in industrial crop Jerusalem artichoke (*Helianthus tuberosus* L.) leaves, *Industrial Crops and Products*, 47: 339-345 (2013).
5. T. YEŞİLOĞLU, B. YILMAZ, B. ÇİMEN, M. İNCESU. Influences of Root stocks on Fruit Quality of 'Henderson' Grapefruit, *Turkish Journal of Agricultural and Natural Sciences*, 1: 1322-1325 (2014).
6. P. TRIPATHI, N.K. DUBEY, A.K. SHUKLA. Use of some essential oils as post harvest botanical fungicides in the management of grey mould of grapes caused by *Botrytis cinerea*, *World J. Microbiol. Biotechnol.*, 24: 39-46 (2008).
7. E.M. SOYLU, F.M. TOK, S. SOYLU, A.D. KAYA, G.A. EVRENDİLEK. Antifungal Activities of the Essential Oils on Post-harvest Disease Agent *Penicillium digitatum*, *Pakistan Journal of Biological Sciences*, 8: 25-29 (2005).
8. A. AMIRI, G. BOMPEIX. Control of *Pencillium expansum* with potassium phosphite and heat treatment, *Crop Protection*, 30: 222-227 (2011).

9. I. TALIBI, L. ASKARNE, H. BOUBAKER, E.H. BOUDYACH, F. MSANDA, B. SAADI, A. AIT BEN AOUMAR. Antifungal activity of some Moroccan plants against *Geotrichum candidum*, the causal agent of post harvest citrus sour rot, *Crop Protection*, 35: 41-46 (2012).
10. A.C. SHUKLA, R.S. YADAV, S.K. SHAHI, A. DIKSHIT. Use of Plant Metabolites as an Effective Source for The Management of Post Harvest Fungal Pests: A review, *Current Discovery*, 1: 33-45 (2012).
11. S.M. AL-REZA, A. RAHMAN, Y. AHMED, S.C. KANG. Inhibition of Plant Pathogens *In vitro* and *In vivo* with Essential Oil and Organic Extract of *Cestrum nocturnum* L., *Pesticide Biochemistry and Physiology*, 96: 86-92 (2010).
12. M.A. PASSONE, N.S. GIRARDI, M. ETCHEVERRY. Antifungal and Antiaflatoxic Activity by Vapor Contact of Three Essential Oils, and Effects of Environmental Factors on Their Efficacy, *LWT-Food Science and Technology*, 53: 434-444 (2013).
13. E.M. SOYLU, Ş. KURT, S. SOYLU. *In vitro* and *in vivo* antifungal activities of the essential oils of various plants against tomato grey mould disease agent *Botrytis cinerea*, *International Journal of Food Microbiology*, 143: 183-189 (2010).
14. M.F. ABDEL-MONAIM, K.A.M. ABO ELYOUSR, K.M. MORSY. Effectiveness of plant extract on suppression of damping off and wilt diseases of lupine (*Lupinus termis* Forsik, *Crop Protection*, 30: 185-191 (2011).
15. M.A. PASSONE, N.S. GIRARDI, C.A. FERRAND, M. ETCHEVERRY. *In vitro* Evaluation of Five Oils as Botanical Fungitoxicants for the Protection of Stored Peanuts from *Aspergillus flavus* and *A. Parasiticus* Contamination, *International Biodeterioration & Biodegradation*, 70: 82-88 (2012).
16. E.L. DE SOUZA, E. DE OLIVEIRA LIMA, K.R. DE LUNA FREIRE, C.P. DE SOUZA. Inhibitory Action of Some Essential Oils and Phytochemicals on the Growth of Various Moulds Isolated From Foods, *Brazilian Archives of Biology and Technology*, 48: 245-250 (2005).
17. CLSI. 2002, Clinical and Laboratory Standarts Institute, formerly NCCLS, National Committee for Clinical and Laboratory Standarts. Reference method for broth dilution antifungal susceptibility testing of yeasts; Approved Standart, 2nd edition, NCCLS document M27-A2, NCCLS, Wayne, PA. Reference method for broth dilution antifungal susceptibility testing of filamentous fungi; Approved Standart, 1st edition, NCCLS document M 38 A, Wayne, PA.
18. L. SVATEZ, F. ZULJAN, M. DERITA, E. PETENATTI, G. TAMAYO, A. CACERES, V.C. FILHO, A. GIMENEZ, R. PINZON, S.A. ZACCHINO, M. GUPTA. Value of the ethnomedical information for the discovery of plants with antifungal properties. A survey among seven Latin American countries, *Journal of Ethnopharmacology*, 127: 137-158 (2010).
19. L. ASKARNE, I. TALIBI, H. BOUBAKER, S.M. AMINE, E.H. BOUDYACH, A. AIT BEN AOUMAR. Effects of organic acids and salts on the development of *Penicillium italicum*: the causal agent of citrus blue mold, *Plant Pathology Journal*, 1: 99-107 (2011).
20. L. ASKARNE, I. TALIBI, H. BOUBAKER, E.H. BOUDYACH, F. MSANDA, B. SAADI, M.A. SERGHINI, A. AIT BEN AOUMAR. *In vitro* and *in vivo* antifungal activity of several Moroccan plants against *Penicillium italicum*, the causal agent of citrus blue mold, *Crop Protection*, 40: 53-58 (2012).
21. V.L. SINGLETON, J.A. ROSSI. Colorimetry of total phenolics with phospho molybdc phosphotungstic acid reagents, *American Journal of Enology and Viticulture*, 16, 144-153 (1965).
22. A. ARNOUS, D.P. MAKKRIS, P. KEFALAS. Effect of principal polyphenolic components in relation to antioxidant characteristics of aged red wines, *Journal of Agricultural and Food Chemistry*, 49: 5736-5742 (2001).
23. N.G. BAYDAR, H. BAYDAR. Phenolic compounds, antiradical activity and antioxidant capacity of oil-bearing rose (*Rosa damascena* Mill.) extract, *Industrial Crops and Products*, 41: 375-380 (2013).
24. Z. ASHRAF, A. MUHAMMAD, M. IMRAN, A.H. TAREQ. *In vitro* Antibacterial and Antifungal Activity of Methanol, Chloroform and Aqueous Extracts of *Origanum vulgare* and Their Comparative Analysis, *International Journal of Organic Chemistry*, 1: 257-261 (2011).
25. V. VOGT, D. CIFUENTE, C. TONN, L. SABINI, S. ROSAS. Antifungal activity *in vitro* and *in vivo* of extracts and lignans isolated from *Larrea divaricata* Cav. against phytopathogenic fungus, *Industrial Crops and Products*, 42: 583-586 (2013).
26. R.P. RASTOGI, B.N. MEHROTRA. Glossary of Indian medicinal plants. New Delhi:National Institute of Science Communication, (2002).
27. T. DERESSA, M. WAKJIRA. Antifungal activity of some invasive alien plant leaf extracts against mango (*Mangifera indica*) anthracnose caused by *Colletotrichum gloeosporioides*, *International Journal of Pest Management*, 61: 99-105 (2015).

28. D.D.C. CARVALHO, E. ALVES, R.B. CAMARGOS, D.F. OLIVEIRA, J.R.S. SCOLFORO, D.A. DE CARVALHO, T.R.S. BATISTA. Plant extracts to control *Alternaria alternata* in Murcott tangor fruits, *Revista Iberoamericana de Micología*, 28: 173-178 (2011).
29. C. GARDUNO-PIZANA, L.L. BARRERA-NECHA, Y.R. GOMEZ. Evaluation of the fungicidal activity of leaves powders and extracts of fifteen Mexican plants against *Fusarium oxysporum* f.sp. *gladioli*, *Plant Pathol.J.*, 9: 103-111 (2010).
30. A. AMSALU, L. FIKIRE, M. DIRIBA. The Antifungal Activity of some medicinal plants against coffee berry disease caused by *Colletotrichum kahawe*, *Int J Agric Res.*, 6: 268-279 (2011).
31. C.N. CUTTER. Antimicrobial effect of herb extracts against *Escherichia coli* O157:H7, *Listeria monocytogenes*, and *Salmonella typhimurium* associated with beef, *Journal of Food Protection*, 63: 601-607 (2000).
32. D. RONGAI, P. PATRIZIO, P. BARBARA, M. FILOMENA. Antifungal activity of some botanical extracts on *Fusarium oxysporum*, *Open Life Sci.*, 10: 409-416 (2015).
33. M. TOSUN, S. ERCİŞLİ, M. ŞENGÜL, H. ÖZER, Antioxidant Properties and Total Phenolic Content of Eight *Salvia* species from Turkey, *Biol. Res.*, 42: 175-181 (2009).
34. I.D. RADOJEVIĆ, M.S. STANKOVIĆ, O.D. STEFANOVIĆ, M.D. TOPUZOVIĆ, L.R. ĆOMIĆ. Antioxidative and antimicrobial properties of different extracts from *Sideritis montana* L., *Romanian Biotechnological Letters*, 17: 7160-7168 (2012).
35. M.S. STANKOVIC, M. TOPUZOVIC, S. SOLUJIC, V. MIHAILOVIC. Antioxidant activity and concentration of phenols and flavonoids in the whole plant and plant parts of *Teucrium chamaerdys* L. var. *glanduliferum* Haussk., *Journal of Medicinal Plants Research*, 4: 2092-2098 (2010).
36. M.S. STANKOVIC, N. NICIFOROVIC, V. MIHAILOVIC, M. TOPUZOVIC, S. SOLUJIC. Antioxidant activity, total phenolic content and flavonoid concentrations of different plant parts of *Teucrium polium* L. subsp. *Polium*, *Acta Societatis Botanicorum Poloniae*, 8: 117-122 (2012).
37. M. TRIGUI, A.B. HSOUNA, S. TOUNSI, S. JAOUA. Chemical composition and evaluation of antioxidant and antimicrobial activities of Tunisian *Thymelaea hirsuta* with special reference to its mode of action, *Industrial Crops and Products*, 41: 150-157 (2013).