

Research on The Detection of Polycyclic Aromatic Hydrocarbons (PAHs) from Fish and Smoked Fish Samples, The Values Obtained and The Significance of Their Presence on Human Health

Received for publication, November , 15, 2016

Accepted, February, 13, 2017

PUCHIANU GHEORGHE¹, DOGARU IULIA³, ENACHE DORIN VALTER¹,
NECULA VALENTIN¹, IONIȚĂ CARMEN²

¹ Transilvania University of Brasov, Faculty of Food and Tourism

² University of Agronomic Sciences and Veterinary Medicine Bucharest

³ Sanitary Veterinary and Food Safety of Brasov

Aderess for correspondence to: gpuchianu@yahoo.com

Abstract

A total of 42 samples of fish and smoked fish products from 22 assortments were analyzed to detect Polycyclic Aromatic Hydrocarbons (PAHs). The sample collection was performed in accordance with the provisions of EC Regulation 333/2007 with its subsequent modifications and additions, and the used detection method was the HPLC (High Performance Liquid Chromatography), validated and accredited by the National Accreditation Body. The obtained values were between undetectable/non-quantifiable and $3.57 \pm 7.67 \mu\text{g}/\text{kg}$ for benzo[a]pyrene and between undetectable/non-quantifiable and $25.65 \pm 10.660.31 \mu\text{g}/\text{kg}$ for the PAH4. The values are below the maximum permitted levels of Regulation (EC) 1881/2006 with its subsequent amendments and additions. Even if the obtained values are within the tolerance limits, fish and smoked fish products must be monitored and examined by laboratory examinations, and the establishment of maximum levels for benzo[a]pyrene and PAH4 are extremely important for protecting the public health. Until now, no data have been submitted in Romania referring to the expertise of fish and smoked fish products for the detection of PAHs.

Keywords: Polycyclic Aromatic Hydrocarbons (PAHs), benzo[a]pyrene, PAH4, expertise

1. Introduction

Polycyclic Aromatic Hydrocarbons (PAHs) are a class of organic compounds containing two or more combined (condensed) aromatic rings. (I.A. DOGARU [1]). They include a number of carcinogenic and genotoxic substances, of which the most important are benzo[a]pyrene, benzo[a]anthracene, benzo[b]fluoranthene, benzo[k]fluoranthene, Benzo[ghi]perylene, Chrysene, Cyclopenta[cd]pyrene, dibenzo[ah]anthracene, dibenzo[ae]pyrene, dibenzo[ai]pyrene, dibenzo[al]pyrene, indeno[1,2,3-cd]pyrene and 5-methylcrisene (G.K. ONTIZAN & al, [5], FOOD STANDARDS AGENCY [14] [15]).

PAHs have a pronounced carcinogenic potential, which has been described since 200 years ago. In 1775, for the first time, English physicist Percival Pott described a correlation between the incidence of scrotum cancer at chimney sweepers and continuous contact with PAH-rich ash (G. PUCHIANU & al, [8]).


The formation of PAH in food depends on the thermal processing methods, correlated with the chemical composition of the product, the way it is packaged, plus the structure of the coating. In this regard, it should be noted that most food preparation processes are performed at temperatures between 370-390 ° C, but also at temperatures higher than 400-600 ° C (eg in the case of fat frying). At temperatures between 400 and 1000 ° C, the PAHs content increases

linearly, and the phenols and fatty acids reach the maximum amount at 600 °C. The smoking temperature between 20 and 55 °C does not significantly affect the production of PAHs, and 40 °C temperature allows for superior flavors than the 20 °C or 55 °C.


In roasted meat (roasting with charcoal grill, frying) PAHs are formed during the incomplete combustion process or by the pyrolysis of the fat, when oxygen and carbon are released, which lead to the formation of polycyclic compounds. These compounds are transported in the resulting smoke and then deposited on meat (A. FARHADIAN & al, [2]). PAHs concentration in meat roasted at a higher distance from coal is lower than in meat roasted closer to coal (M.D. GUILLEN & al, [4]). In order to avoid the formation of large amounts of PAHs, grilled meat should not be in contact with the flame, the process should last longer and the fat should be kept as low as possible (W. STALL & al, [9], S. TAO & al, [10]).

Another PAHs-generating process is smoking, a type of food preservation, which has been correlated with the increased incidence of pharyngeal cancer, observed in Icelandic fishermen, large smoked fish consumers, even if PAHs were detected in low concentrations. Accumulation of PAHs in smoked food depends on how food was smoked, the type of the fuel, the temperature and duration of smoking. Wood quality plays an important role, so pyrolysis of hardwood generates poorer smoke in PAHs than soft woods that contains resins (G.. GRIMMER [3]). PAHs are mainly formed from the carbohydrates found in food, at high temperatures and in the absence of oxygen.


As a contaminant, the marker is considered benzo(a)pyrene and the PAH4: benzo(a)pyrene (C₂₀H₁₂), benzo(a)anthracene (C₁₈H₁₂), benzo(b)fluoranthene (C₂₀H₁₂) and chrysene (C₁₈H₁₂) ([http://ro.wikipedia.org/wiki/, Benzo \[a\] piren \[22\]](http://ro.wikipedia.org/wiki/Benzo_a_piren), [http://ro.wikipedia.org/wiki/, Benz \[a\] antracen \[23\]](http://ro.wikipedia.org/wiki/Benz_a_antracen), [http://ro.wikipedia.org/wiki/, Benzo \[b\] fluoranten \[24\]](http://ro.wikipedia.org/wiki/Benzo_b_fluoranten), [http://ro.wikipedia.org/wiki/, Crisen \[25\]](http://ro.wikipedia.org/wiki/Crisen)).




C₂₀H₁₂



C₁₈H₁₂



C₂₀H₁₂



C₁₈H₁₂

Depending on the number of rings, hydrocarbons can be divided into:

- light PAHs (2-4 aromatic rings): chrysene; benz(a)anthracene;
- heavy PAHs (with 5 or more aromatic rings): benzo(a)pyrene; benzo(a)fluoranthene; dibenzo(a)anthracene.

Benzo[a]pyrene is a polycyclic aromatic hydrocarbon found in coal tar. Its metabolites are mutagenic and highly carcinogenic (classified as a Carcinogenic Group 1 by IARC), being a result of the transformation of starch, amino acids and fatty acids during incomplete combustion at temperatures between 300-600 °C, following. (IARC [17]). The content is expressed in micrograms / kg ($\mu\text{g} / \text{Kg} = \text{ppb}$). The carcinogenic action of benzoprene can be obtained by direct contact (dermal), and especially by ingestion (A.R. NAGHIU & al, [7]).

2. Materials and Methods

Within the Sanitary Veterinary and Food Safety Laboratory in Brasov, a number of 42 samples of fish and smoked fish products from 22 assortments were analyzed for the detection of Polycyclic Aromatic Hydrocarbons (SPECIFIC PROCEDURE [20]).

Sampling was carried out in accordance with the provisions of EC Regulation 333/2007 which determines the sampling methods and methods of analysis for the official control of the levels of Pb, Cd, Hg, 3-MCPD and PAHs in food, with subsequent amendments and additions (EU REGULATION no. 333/2007 [13]).

Thus, sampling was carried out by trained persons, with due regard for measures designed to eliminate the risk of accidental contamination of the prelevated samples and to avoid any changes that could have affected contaminant levels, which would have had a negative effect on analytical determination or who would have made the evidences unrepresentative.

The overall sample was about 1 kg, unless this was not possible, for example, when the sample consisted of a package or a unit.

The minimum number of incremental samples taken from the lot or subplot was established as follows: (Table 1).

Table 1. The minimum number of incremental samples to be taken from the batch or subplot

Lot / subplot weight or volume (in kg or liter)	Minimum incremental number of samples to be sampled
< 50	3
≥ 50 și ≤ 500	5
> 500	10

If the batch or subplot was made up of separate packages or units, the number of packages or units to be sampled to form a bulk was determined according to table 2.

Table 2. Number of packages or units (incremental samples) to be taken to constitute the bulk sample if the consignment or subplot was made up of separate packages or units

Number of packages or units in the lot/sublot	Number of packages or units to be taken
≤ 25	at least 1 pack or 1 unit
26-100	about 5%, at least 2 packages or units
> 100	about 5%, up to 10 packages or units

After sampling, each sample was placed in a clean and inert container (aluminum foil) which provides adequate protection against contamination or adsorption of substances on the inside of the container and damage during transport. The taken samples were sealed and identified, keeping records of each sampling.

The detection method used was high performance liquid chromatography (HPLC), a qualitative analysis method used in biochemistry and analytical chemistry for the separation, identification and quantification of compounds, validated and accredited by the National Accreditation Body (RENAR) (SR EN ISO 15753 / 2007 [21], A.D. NAGHIU & al, [6], M.A. QUILLIAM [11]).

The principle of the method is that PAHs are extracted in an acetonitrile/acetone mixture followed by two purifications on the C18 reverse phase cartridge and on the florisil cartridge. After separation, liquid chromatography is determined and fluorescence measured at different excitation and emission wavelengths, according to the standardized and optimized working method. In order to obtain repeatable results, the ambient temperature in the

laboratory was stable ($\leq 20^\circ\text{C}$), higher temperatures, increasing the solubility of short chain fatty acids. The detection program is presented in Table 3.

Table 3. Detection program

Component	Duration min	Excitation wavelength nm	Emission wavelength nm
<i>Benz[a]anthracene</i>	12.5	270	385
<i>Chrysene</i>	14.0		
<i>Benzo[b]fluoranthene</i>	21.8	256	446
<i>Benzo[a]pyrene</i>	31,2	292	410
<i>Dibenz[a,h]anthracene</i>	44.1		

After weighing exactly 2 gramms of the sample, 5 steps(0 - 4) were carried out for the analytical process, followed by the actual determination (optimization of the apparatus, calibration of the calibration curve, reading, calculation and expression of the results).

Step 0

The sample to be analyzed, chopped and well homogenized, was extracted with methanol and then with chloroform, in two steps. The last bottom layer was collected in the same vial and then evaporated to dryness (Figures 1 and 2).




Fig.1. Extraction of solid samples



Fig.2. Collecting lowest layer

Thus, the fat was obtained

Step 1

The sample to be analyzed was subjected to a three-step extraction with 'Mixture1', the supernatant (3 x 4 ml) being collected in the same vial, then evaporated to dryness. The fatty residue was thus obtained (Figure 3).




Fig.3. Samples after evaporation to dryness – fatty residue

Step 2

The fat residue was purified by successive extractions with the same 'Mixture1' in three steps, the supernatant (3 x 2 ml) being collected in the same vial, and then purified on the C18 cartridge. Prior to passing through the cartridge, it was conditioned according to the manufacturer's operating instructions (Figure 4).




Fig. 4. Conditioning the cartridge

After passing the eluate, 'Mixture 1' was added to elute all the particles of interest. All the eluate thus obtained was evaporated to dryness. The resulting oily residue was treated with hexane and thus the hexane extract was obtained, which was kept for 24 hours at -18 °C.

Step 3

The hexane extract was subjected to successive extractions with 'Mixture 3' in three steps, the supernatant (3 x 2 ml) was collected in the same vial and then purified on the florisil cartridge (Figure 5)



Fig 5. Purification of the supernatant through the florisil cartridge

Prior to passing through the florisil cartridge, it was conditioned according to the manufacturer's instructions. After passing the eluate, Mixture 3 was added to elute all the particles of interest.

Step 4

All the eluate thus obtained was concentrated to 1 ml. It was then treated with 0.5 ml of toluene and evaporated to 50 µl, internal standard was added and completed up to the volume of interest (250, 500 µl) with acetonitrile. Note that all of the fat residues were weighed and the evaporation to dryness was done with nitrogen in a water bath at 350 °C.

Determination Steps:

Optimizing the device

The machine was started before the determination and it started the optimization of the working parameters. (Figure 6)



Fig. 6. Varian LC MS

Marking the standard curve

PAHs were prepared at concentrations of 1, 2, 3, 4, 5 ppb, with which the calibration curve was drawn. The curve shall be linear and pass through the origin, and the correction coefficient shall be at least 0.995.

The readout

After adjusting the device, the identification data of the samples to be analyzed was entered into the program. (Figure 7)




Fig. 7. Inserting sample identification data in the software

After the device was initialized, the samples were injected, following all the steps required by the machine's reading program. The reading was completed by the individual chromatogram for each sample and by an analysis report, which identifies the concentration of each substance at the retention times specific to the analyzed PAHs.

To the determination set it was also added a fortified sample at the maximum allowed according to the legislation in force (EC Regulation 1881/2006, with subsequent amendments and completions) (EU REGULATION no. 1881/2006 [12]).

Calculation and results analysis

Before the calculation, the following aspects were followed: the recovery percentage of the fortified sample to be in accordance with the regulations in force (EC Regulation 333/2007 with subsequent amendments and completions); retention times are appropriate for PAHs analyzed; the internal standard area is the appropriate one; in the case of dilutions, take account of the dilution factor; issuing the analysis report and processing the data by correcting the value with the recovery ratio for Benzo(a)pyrene and the PAH4 respectively.

3. Results and discussion

Following the laboratory examination of 42 samples of fish and smoked fish products, the following results were obtained: (Table 4.)

Table 4. The type of analyzed samples, their number and the obtained results.

Sample	Number of probes	Results		
		benzo[a]pyrene $\mu\text{g}/\text{kg}$	benzo[a]pyrene recuperation %	PAH4 $\mu\text{g}/\text{kg}$
Mackerel pastrami	1	Undetectable*	-	3,24 \pm 1,35
Smoked mackerel	5	Unquantifiable**	-	2,83 \pm 1,18
		Unquantifiable**	-	1,87 \pm 0,78
		Unquantifiable**	-	1,99 \pm 0,83
		Undetectable	-	3,92 \pm 1,53
		Undetectable	-	Unquantifiable**
Smoked mackerel fillet	2	2,29 \pm 0,91	84,9	3,09 \pm 1,28
		2,11 \pm 1,23	86,8	1,98 \pm 0,86
Smoked Trout	3	Unquantifiable**	-	2,99 \pm 1,24
		0,75 \pm 0,30	88,3	7,61 \pm 3,16
		Unquantifiable**	-	1,99 \pm 0,83
Smoked Sprat	3	Unquantifiable**	-	6,30 \pm 2,62
		0,87 \pm 0,35	88,3	7,64 \pm 3,17
		1,37 \pm 0,4306	86,8	11,7813
Smoked Sprat in Sunflower Oil	7	Unquantifiable**	-	Undetectable*
		Undetectable*	-	2,27 \pm 0,94
		Unquantifiable**	-	6,54 \pm 2,72
		Undetectable*	-	0,25 \pm 0,10
		Unquantifiable**	-	2,23 \pm 0,93
		Unquantifiable**	-	2,11 \pm 0,96
		Undetectable*	-	3,67 \pm 1,52
Smoked Sprat in Rapeseed Oil	1	Undetectable*	-	2,57 \pm 1,07
Sprat smoked with beech and alder smoke	2	Undetectable*	-	18,46 \pm 7,67
		Undetectable*	-	4,96 \pm 2,06
Frozen Smoked Sprat	1	3,57 \pm 7,67	-	25,65 \pm 10,66
Mussels in sunflower oil	2	0,76 \pm 0,30	98,32	8,17 \pm 3,39
		0,84 \pm 0,33	98,32	9,73 \pm 4,04
Smoked Herring	1	Unquantifiable**	-	9,88 \pm 4,10
Sliced smoked salmon	1	Unquantifiable**	-	2,32 \pm 0,96
Smoked salmon filet	2	Undetectable*	-	1,39 \pm 0,58
		Undetectable*	-	1,39 \pm 0,58
Smoked Shells in oil	2	Unquantifiable**	98,32	7,84 \pm 3,26
Smoked Bighead Filet	1	Undetectable*	-	0,80 \pm 0,33
Vaccumed Smoked	1	Undetectable*	-	0,69 \pm 0,29

Tuna Filet				
Smoked Pontic Shad	2	Unquantifiable**	-	2,3128 ±1,3749
		Unquantifiable**	-	2,32 ±0,96
Smoked Herring Filet	1	Undetectable*	-	1,72±0,71
Doripesco Fish Rod	1	Unquantifiable**	-	1,92±0,61
Frozen mussels pasteurized in vacuum	1	Undetectable*	98,32	2,89 ±1,20
Mussels in smoke flavored oil	1	1,38±0,55	69,4	6,94±2,88
Smoked Mussels in Sunflower Oil 'Franz'	1	0,79±0,31	-	5,54±2,30

* Undetectable < LOD (Limit of detection, smallest measured content, from which it is possible to deduce the presence of the analyte with reasonable statistical certainty.)

** unquantifiable < LOQ (Limit of quantification, lowest content of the analyte which can be measured with reasonable statistical certainty.)

The obtained values were between undetectable/non-quantifiable and $3.57 \pm 7.67 \mu\text{g} / \text{kg}$ for Benzo(a)pyrene and between undetectable/non-quantifiable and $25.65 \pm 10,660.31 \mu\text{g} / \text{kg}$ for PAH4 (corrected with recovery). All these values are below the maximum permitted levels specified in Regulation (EC) 1881/2006 with subsequent amendments and additions. (Table 5)

Table 5. Maximum allowable limits for PAHs in fishmeal and smoked fish products (EU REGULATION no. 1881/2006 [12])

Maximum limits ($\mu\text{g}/\text{kg}$)	
Benzo[a]pyrene	Sum of Benzo[a]pyrene, Benz[a]anthracene, Benzo[b]fluoranthene and Chrysene
5.0 $\mu\text{g}/\text{kg}$ before 1.09.2014 2.0 $\mu\text{g}/\text{kg}$ after 1.09. 2014	30.0 $\mu\text{g}/\text{kg}$ before 1.09.2014 12.0 $\mu\text{g}/\text{kg}$ after 1.09.2014

There is an exception for smoked sprats and canned smoked sprats (*sprattus sprattus*); bivalve molluscs (fresh, chilled or frozen); heat treated meat and heat treated meat products sold to the final consumer, where the admissibility limit for Benzo[a]pyrene is $5.0 \mu\text{g}/\text{kg}$, and for PAH5 is $35.0 \mu\text{g}/\text{kg}$. Also for bivalve mollusks, the admissibility limit is $6.0 \mu\text{g}/\text{kg}$, and for PAH4 is $35.0 \mu\text{g}/\text{kg}$. (FOOD SAFETY AUTHORITY OF IRELAND [16])

The highest value for Benzo[a]pyrene was detected in the Frozen Smoked Sprat samples, $3,57 \pm 7,67 \mu\text{g}/\text{kg}$. It is higher than the maximum admissible value ($5,0 \mu\text{g}/\text{kg}$) with $1,43 \mu\text{g}/\text{kg}$. Also high values were detected in the Smoked mackerel samples ($2,29 \pm 0,91 \mu\text{g}/\text{kg}$), in the Mussels in smoke flavored oil samples ($1,38 \pm 0,55 \mu\text{g}/\text{kg}$) and in the Smoked Sprat samples ($1,37 \pm 0,4306 \mu\text{g}/\text{kg}$).

As for PAH4, the highest value was detected in the Frozen Smoked Sprat samples ($25,65 \pm 10,6667 \mu\text{g/kg}$), higher than the maximum admissible value of $35,0 \mu\text{g/kg}$, resulting a difference of $9,35 \mu\text{g/kg}$. High values were also detected in the Sprat smoked with beech and alder smoke samples ($18,46 \pm 7,67 \mu\text{g/kg}$), in the smoked herring samples ($9,88 \pm 4,10 \mu\text{g/kg}$), in the Mussels in sunflower oil samples ($9,73 \pm 4,04 \mu\text{g/kg}$, respectively $8,17 \pm 3,39 \mu\text{g/kg}$) and in the Smoked Shells in oil samples ($7,84 \pm 3,26 \mu\text{g/kg}$).

The obtained values from the other food products were lower for both parameters, the lowest values being recorded for benzo(a)pyrene in the smoked mackerel samples, smoked trout, Smoked Sprat in Sunflower Oil, and in the herring, salmon, tuna, shells specialities, and, in case of PAH4, in the smoked mackerel and Smoked Sprat in Sunflower Oil samples.

Even if the obtained values are within the admissible limits, fish and smoked fish products must be monitored and tested by laboratory examinations, and setting the maximum levels for benzo(a)pyrene and for PAH4 are very important since they are meant to protect the public health.

This is why, annually, the National Sanitary Veterinary and Food Safety Authority establishes a sampling plan for both animal and non-animal food samples in order to monitor and detect PAHs, for the purpose of preventing possible risks as follows: (Table 6)

Table 6. Control of contaminants in food of animal origin
(ORDER OF THE N.S.V.F.S.A. 35/2016 [19], and 51/2005 [18])

Substance	Matrix	Prelevation source	Analysis method	
Benzopyrene	Smoked meat and smoked meat products	Processing units, marketing units	Screening	Confirmation

In case of smoked fish and fish products, maintaining levels of PAHs within acceptability limits can reasonably be achieved by observing good manufacturing practices and by applying prevention measures against food contamination.

In order to ensure effective public health protection, fish and smoked fish products containing excess PAHs must not be placed on the market as such or mixed with other foods nor used as ingredients.

4. Conclusion

The maximum values of PAHs were obtained from the examination of frozen smoked sprat. Thus in the case of benzo(a)pyrene the value was $3.57 \pm 7.67 \mu\text{g/kg}$, and in the case of PAH4 the value was $25.65 \pm 10.6667 \mu\text{g/kg}$. The values are within the admissibility limits, but they are close to the maximum admissible limits.

Higher values, but within tolerance limits, were obtained for smoked mackerel for benzo(a)pyrene and for sprat smoked in beech and alder smoke in the case of PAH4.

The lowest values were recorded for benzo(a)pyrene, in smoked mackerel, smoked trout, smoked sprat in oil and herring, mackerel, salmon, tuna, novak, shells samples., and in the case of PAH4, for smoked mackerel and smoked sprat in oil samples.

The obtained results show that fish and smoked fish products have not been a hazard to the human consumer, but values close to the maximum allowable values are an argument for the requirement of smoked food to be tested for PAHs in order to prevent the illness of consumers.

Until now, no data has been presented in Romania referring to the evaluation of fish and smoked fish products in regard to PAHs values.

5. Competing interest

The authors declare no conflict of interest.

References

1. DOGARU IULIA ANNABELLA. 2014. Research on the significance of PAH presence in different foods and the results of their detection. Dissertation paper - Transilvania University of Brasov, (2014).
2. FARHADIAN, A., JINAP, S. FARIDAH A., SAKAR, Z.I. Determination of Polycyclic aromatic hydrocarbons in grilled meat, *Food Control*, Volume 21, (2010).
3. GRIMMER, G. Environmental Carcinogens: Polycyclic aromatic hydrocarbons, Boca Raton, Florida, CRC Press (2017).
4. GUILLEN, MD., SOPELANA, P., AND PARTEARROYO, M.A. Food as a source of polycyclic aromatic carcinogens, *Rev. Environ. Health*, (1997).
5. MONTIZAAN, G.K., KARMERS., P.G.N., JANUS, J.A., SI POSTHUMUS, R. Integrated Criteria Document Polynuclear aromatic hydrocarbons (PAH): Effects of 10 selected compounds. Appendix to RIVM Report no. 758474007, National Institute of Public Health and Environmental Protection, RIVM, Bilthoven, (1989).
6. NAGHIU, A.,D. SIMEDRU, C. LASLO, A. MIHALTAN. An easy and reliable method for PAH extraction from Food samples, *Agricultura, agricultural practice and science journal*, Vol 79, nr.1-2, (2012).
7. NAGHIU, A., R. LASLO, D. SIMEDRU, C. LASLO. The present of PAH's in meat products found on the Cluj-Napoca market, *Agricultura, agricultural practice and science journal*, Vol 79, nr.1-2, (2012).
8. PUCHIANU GHEORGHE, DOGARU IULIA ANNABELLA, MĂRCULESCU ANGELA. Research on detection of PAHs in samples of meat and smoked meat products, the values obtained, their presence and significance on human health. 3rd North and East European NEEFOOD Congress on Food. Global and Local Challenges in Food Science and Technology, Braşov, România, (2015).
9. STALL, W., AND EISENDRAND, G. Determination of Polynuclear Aromatic Hydrocarbons and Nitrosamines, HPLC in *Food Analysis, Academic Press, New York*, (1998).
10. TAO S., and COL. Dispersion modeling of Polycyclic aromatic Hydrocarbons from combustion of biomass and fossil fuels and production of coke in Tianjin, *Environ Sci Technol* 40, (2006).
11. QUILLIAM M.A. and SIM P.G. Determination of polycyclic aromatic compounds by high-performance liquid chromatography with simultaneous mass spectrometry and ultraviolet diode array detection, *J Chromatogr Sci*, 26, (1988).
12. EU REGULATION no. 1881/2006: Setting the maximum level for certain contaminants in foodstuffs, with subsequent amendments, (2006).
13. EU REGULATION no. 333/2007: Establishment of sampling methods and methods of analysis for the official control of Pb, Cd, Hg, 3-MCPD, and PAH content in foodstuffs, with subsequent amendments and supplemented, (2007)
14. FOOD STANDARDS AGENCY. PAHs in dietary supplements, *Food Survey Information Sheet* 86/05, (2005).
15. FOOD STANDARDS AGENCY. Polycyclic aromatic hydrocarbons (PAHs) in baby foods and infant formulae, *Food Survey Information Sheet*, (2006).
16. FOOD SAFETY AUTHORITY OF IRELAND. Polycyclic Aromatic Hydrocarbons (PAHs) in Food, Issue no.2/June 2, (2015).
17. IARC. Certain Polycyclic Aromatic Hydrocarbons and Heterocyclic Compounds, IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemical to Humans, Vol 3, *International Agency for Research on Cancer, Lyon*, (1973).
18. ORDER OF THE NATIONAL SANITARY VETERINARY AND FOOD SAFETY AUTHORITY, no. 51/2005: Sanitary veterinary norms for the implementation of the measures for supervision and control of some substances and their residues in live animals and products through the performance of the analytical methods and interpretation of the results.
19. ORDER OF THE NATIONAL SANITARY VETERINARY AND FOOD SAFETY AUTHORITY, no. 35/2016. Methodological Norms for the Application of the Action Program for the Control, Prevention, Control and Eradication of Animal Diseases, Transmissible Animal Diseases, Animal Protection and Methodological Norms for the Application of the Food Safety Supervision and Control Program.

20. Specific Procedure (PS-13-DTSAONASRA) – The National Sanitary Veterinary and Food Safety Authority - Specific procedure for sampling non-animal products.
21. SR EN ISO 15753 / 2007- animal and vegetable fats and oils. Determination of PAHs.
22. <http://ro.wikipedia.org/wiki/>, Benzo [a] piren.
23. <http://ro.wikipedia.org/wiki/>, Benz [a] antracen.
24. <http://ro.wikipedia.org/wiki/>, Benzo [b] fluoranten.
25. <http://ro.wikipedia.org/wiki/>, Crisen.